



**University of  
Zurich**<sup>UZH</sup>

**Zurich Open Repository and  
Archive**

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2012

---

## **Chiral Imidazoles and Imidazole N-Oxides as Ligands for Stereoselective Cyclopropanation reactions**

Mlostoń, G ; Mucha, P ; Heimgartner, H

Abstract: New, optically active ligands derived from imidazole were tested in the cyclopropanation reaction of styrene using ethyl diazoacetate and copper(I) triflate as a catalyst.

DOI: <https://doi.org/10.2174/157017812800221735>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-61481>

Journal Article

Accepted Version

Originally published at:

Mlostoń, G; Mucha, P; Heimgartner, H (2012). Chiral Imidazoles and Imidazole N-Oxides as Ligands for Stereoselective Cyclopropanation reactions. *Letters in Organic Chemistry*, 9(2):89-91.

DOI: <https://doi.org/10.2174/157017812800221735>

## **Chiral Imidazoles and Imidazole *N*-Oxides as Ligands for Stereoselective Cyclopropanation Reactions**

Grzegorz Mloston<sup>\*a</sup>, Paulina Mucha<sup>a,[1]</sup> and Heinz Heimgartner<sup>\*b</sup>

<sup>a</sup> University of Łódź, Department of Organic and Applied Chemistry, Tamka 12, PL-91-403 Łódź, Poland

<sup>b</sup> University of Zurich, Institute of Organic Chemistry, Winterthurerstrasse 190, CH-8057 Zurich, Switzerland

**Abstract:** New, optically active ligands derived from imidazole were tested in the cyclopropanation reaction of styrene using ethyl diazoacetate and copper(I) triflate as a catalyst.

**Key words:** cyclopropanation, diazo compounds, styrene, imidazole *N*-oxides

**Running header:** Bis-imidazole derivatives as chiral ligands in cyclopropanation reactions

---

<sup>\*</sup> Corresponding authors: G. M., E-mail: gmloston@uni.lodz.pl, Tel: +48 42 635 57 61, Fax: +48 42 665 51 62; H. H., E-mail: heimgart@oci.uzh.ch, Tel: +41 44 635 42 82, Fax: +41 44 635 68 12

## Introduction


Cyclopropanation reactions are well known and widely applied in modern organic synthesis [2]. On the one hand, diverse cyclopropane derivatives are important building blocks for the preparation of more complex molecules, e.g. natural products [3], and on the other hand, cyclopropanations are frequently used as test reactions in basic research. Special attention is focused on stereoselective formation of the cyclopropane ring, and both diastereoselectivity and enantioselectivity are of interest [4]. A favorite procedure for the examination of new chiral ligands is the metal complex catalyzed cyclopropanation of prochiral olefins with  $\alpha$ -diazo compounds. The most popular substrates for these reactions are styrene and its derivatives and  $\alpha$ -diazo esters [4e].

In recent times, chiral *N*-heterocycles and their *N*-oxides have attracted attention as ligands for asymmetric synthesis [5]. However, there are only a few reports on the application of imidazoles and imidazole *N*-oxides [6]. In a series of recent papers, we reported on the synthesis of differently substituted, optically active, 2-unsubstituted imidazole 3-oxides and their conversion into other imidazole derivatives, e.g. imidazole-2-thiones [7]. To the best of our knowledge, neither imidazoles nor their *N*-oxides have been used in asymmetric cyclopropanation.

The goal of the present study was to test selected bis-imidazoles and the corresponding *N,N'*-dioxides, derived from enantiomerically pure *trans*-1,2-diaminocyclohexane, as ligands in the Cu(I)-catalyzed reaction of styrene with ethyl diazoacetate (EDA).


## Results and discussion

The preparation of the chiral ligands **1-4** (**Fig. (1)**) has been reported previously [7].



**Figure 1.** Imidazole derived ligands **1-4** used in the study

The first cyclopropanation reaction, in which styrene and EDA in a ratio of 4:1 were reacted in the presence of Cu(I) triflate as a catalyst, was carried out at room temperature without the addition of a ligand. The progress of the reaction was monitored by IR spectroscopy and the disappearance of the strong diazo absorption at  $2111\text{cm}^{-1}$  evidenced the completion of the reaction. The mixture of *trans*- and *cis*-2-phenylcyclopropyl carboxylate **5** was isolated in 60% yield, and the ratio of the *trans*- and *cis*-isomer was established by  $^1\text{H}$ -NMR spectroscopy as 2:1 (Scheme 1) (see *e.g.* [8]).



**Scheme 1.** Cyclopropanation of styrene with ethyl diazoacetate (EDA) in the presence of CuOTf.

An analogous experiment performed in the presence of catalytic amounts (4 mol % relative to EDA used) of the bis-imidazole 3,3'-dioxide  $(S,S)$ -**1a** led to a complex mixture of products. The reaction to give the desired cyclopropane derivatives occurred smoothly only in the presence of trace amounts of hydrazine hydrate [9][10]. After chromatographic workup, the determination of the *ee*-values for *trans*- and *cis*-**5** was carried out by HPLC on a chiral solid phase (Chiralcel OD). This method was applied for all studied ligands (Table 1).

In general, the diastereoisomeric excess (*de*) observed in the reactions performed in the presence of the chiral ligands **1-4** was slightly higher than in their absence.

**Table 1.** Efficiency of imidazole ligands **1-4** in the cyclopropanation reaction of styrene using ethyl diazoacetate

Ligand	Yield [%]	<i>trans</i> - <b>5</b> : <i>cis</i> - <b>5</b>	<i>ee trans</i> - <b>5</b> [%]	<i>ee cis</i> - <b>5</b> [%]
( <i>S,S</i> )- <b>1a</b>	52	3:1	-	-
( <i>S,S</i> )- <b>1b</b>	34	2.3:1	4	31
( <i>S,S</i> )- <b>1c</b>	60	4.4:1	2	34
( <i>S,S</i> )- <b>1d</b>	50	2:1	20	40
( <i>R,R</i> )- <b>2b</b>	48	2.5:1	14	94
( <i>S,S</i> )- <b>2c</b>	29	2.3:1	8	34
( <i>R</i> )- <b>3</b>	41	3:1	-	-
( <i>S</i> )- <b>4</b>	54	3:1	1	4

The highest *de* value (81%) was obtained with (*S,S*)-**1c**. The determined *ee*-values were rather low, but they were higher in the case of *cis*-**5** than in *trans*-**5**. The best induction for *cis*-**5** (94% *ee*) was achieved with the ligand (*R,R*)-**2b**. On the other hand, the highest *ee* of *trans*-**5** (20%) was observed in the reaction with (*S,S*)-**1d**.

Mixtures of isomeric cyclopropanecarboxylates **5** obtained in the presence of catalytic amounts of catalysts (*S,S*)-**1d** and (*R,R*)-**2b**, respectively, were separated and fractions containing analytically pure *trans*-**5** were obtained. In both cases, the optical rotatory power was measured. Whereas in the first case the  $[\alpha]_D$  value was +57 (*c* 0.2; CHCl<sub>3</sub>), in the second case it was -31 (*c* 0.2; CHCl<sub>3</sub>). Based on the literature data for the optical rotation of (1*S*,2*S*)-*trans*-**5** ( $[\alpha]_D = +269$ ) and (1*R*,2*R*)-*trans*-**5** ( $[\alpha]_D = -279$ ) [12], one can conclude that in the experiments with (*S,S*)-**1d** the first enantiomer predominated and, in the case of (*R,R*)-**2b** used as a catalyst, the second enantiomer was the major component.

## Conclusions

The cyclopropanation reaction of styrene with EDA in the presence of CuOTf and catalytic amounts of hydrazine was studied in order to test the efficiency of chiral

imidazole derivatives as ligands with respect to the stereoselectivity of the process. In comparison with the ‘ligand-free’ reaction, the *de*-value in favour of *trans*-**5** in general increased slightly, and in the case of (*S,S*)-**1c** achieved the value of 81%. The influence of the ligands **1-4** on the *ee*-value was modest and only in the case of (*R,R*)-**2b** was an attractive result (94%) for the *cis*-isomer obtained. It is very likely that the observed chiral induction in reactions catalyzed by enantiopure ligands derived from imidazole *N*-oxides and imidazole derivatives results from complexation of the Cu(I) salt and formation of a chiral complex, which is incorporated in the structure of the intermediate carbenoid. The structures of some chiral carbenoids have been proposed for metal catalyzed cyclopropanation reactions with diazoacetates and styrene [4e]. To the best of our knowledge, the presented ligands are the first imidazole derivatives tested in the cyclopropanation reaction. In spite of the relatively modest selectivities observed, the presented results are encouraging for further modifications of the substitution pattern in chiral imidazole derivatives.

## References

- [1] Part of the Ph. D. Thesis of *P. M.*, University of Łódź, 2010.
- [2] a) Davies, H. M. L.; Antoulinakis, E. G. Intermolecular Metal-Catalyzed Cyclopropanations. *Org. React.*, **2001**, 57, 1–326; b) Charette, A. B.; Beauchemin, A. Simmons-Smith Cyclopropanation Reaction. *Org. React.*, **2001**, 58, 1–415.
- [3] Wessjohann, L. A.; Brandt, W.; Thiemann, T. Biosynthesis and Metabolism of Cyclopropane Rings in Natural Compounds. *Chem. Rev.*, **2003**, 103, 1625–1647.
- [4] a) Lebel, H.; Marcoux, J.-F.; Molinaro, C.; Charette, A. B. Stereoselective Cyclopropanation Reactions. *Chem. Rev.*, **2003**, 103, 977–1050; b) Maas, G. Ruthenium-catalysed Carbene Cyclopropanation Reactions with Diazo Compounds. *Chem. Soc. Rev.*, **2004**, 33, 183–190; c) Kanchiku, S.; Suematsu, H.; Matsumoto, K.; Uchida, T.; Katsuki, T. Construction of an Aryliridium-Salen Complex for Highly *cis*- and Enantioselective Cyclopropanations. *Angew. Chem. Int. Ed.*, **2007**, 46, 3889–3891; d) Ichinose, M.; Suematsu, H.; Katsuki, T. Efficient Construction of  $\alpha$ -Spirocyclopropyl Lactones: Iridium-Salen-Catalyzed Asymmetric Cyclopropanation. *Angew. Chem. Int.*

*Ed.*, **2009**, 48, 3121–3123; e) Caballero, A., Prieto, A., Diaz-Requejo, M. M., Perez, P. J., Metal-Catalyzed Olefin Cyclopropanation with Ethyl Diazoacetate: Control of the Diastereoselectivity. *Eur. J. Org. Chem.*, **2009**, 1137–1144.

[5] a) Malkov, A. V.; Kočovský, P. Chiral *N*-Oxides in Asymmetric Catalysis. *Eur. J. Org. Chem.*, **2007**, 29–36; b) Kadlčíková, A.; Hrdina, R.; Valterová, I.; Katora, M. Simple and Fast Synthesis of New Axially Chiral Bipyridine *N,N'*-Dioxides for Highly Enantioselective Allylation of Aldehydes. *Adv. Synth. Catal.*, **2009**, 351, 1279–1283; c) Malkov, A. V.; Ramirez-López, P.; Biedermannova, L.; Rulišek, L.; Dufkova, L.; Katora, M.; Zhu, F.; Kočovský, P. On the Mechanism of Asymmetric Allylation of Aldehydes with Allyltrichlorosilanes Catalyzed by QUINOX, a Chiral Isoquinoline *N*-oxide. *J. Am. Chem. Soc.*, **2008**, 130, 5341–5348.

[6] a) Kulhánek, J.; Bureš, F.; Šimon, P.; Schweizer, W. B. Utilizing Terpene Derivatives in the Synthesis of Annulated Terpene-Imidazoles with Application in the Nitroaldol Reaction. *Tetrahedron: Asymmetry*, **2008**, 19, 2462–2469; b) Bureš, F.; Kulhánek, J.; Ružička, A. Probing Electronic and Regioisomeric Control in an Asymmetric Henry Reaction Catalyzed by Camphor-imidazoline Ligands. *Tetrahedron Lett.*, **2009**, 50, 3042–3045; c) Kwiatkowski, P.; Mucha, P.; Mlostoń, G.; Jurczak, J. Novel Chiral *C*<sub>2</sub>-Symmetric Bis-imidazole-*N*-oxides as Promising Organocatalyst for Enantioselective Allylation of Aromatic Aldehydes. *Synlett*, **2009**, 1757–1760.

[7] a) Jasiński, M.; Mlostoń, G.; Mucha, P.; Linden, A.; Heimgartner, H. Synthesis of New Bis-imidazole Derivatives. *Helv. Chim. Acta*, **2007**, 90, 1765–1780; b) Mucha, P.; Mlostoń, G.; Jasiński, M.; Linden, A.; Heimgartner, H. A New Approach to Enantiomerically Pure Bis-imidazoles Derived from *trans*-1,2-Diaminocyclohexane. *Tetrahedron: Asymmetry*, **2008**, 19, 1600–1607; c) Jasiński, M.; Mlostoń, G.; Linden, A.; Heimgartner, H. Synthesis and Selected Transformations of 1*H*-Imidazole 3-Oxides Derived from Amino Acid Esters. *Helv. Chim. Acta*, **2008**, 91, 1916–1933; d) Mlostoń, G.; Jasiński, M. Synthesis and Selected Transformations of 3-Oxido-1*H*-imidazole-4-carboxamides. *Collect. Czech. Chem. Commun.*, **2010**, 75, 871–885.

[8] a) Shepperson, I.; Quici, S.; Pozzi, G.; Nicoletti, M.; O'Hagan, D. *C*<sub>2</sub>-Symmetric Fluorous Diamines and Diimines as Ligands for Metal-Catalysed Asymmetric

Cyclopropanation of Styrene. *Eur. J. Org. Chem.*, **2004**, 4545–4551; b) Le Maux, P.; Abrunhosa, I.; Berchel, M.; Simonneaux, G.; Guela, M.; Masson, S. Application of Chiral 2,6-Bis(thiazoliny)pyridines in Asymmetric Ru-catalyzed Cyclopropanation with Diazoesters. *Tetrahedron: Asymmetry*, **2004**, *15*, 2569–2573.

[9] Hydrazine and its derivatives prevent oxidation of the Cu(I) salt and they are recommended to improve the yields of cyclopropanecarboxylates; see: a) Leutenegger, U.; Umbricht, G.; Fahrni, C.; von Matt, P.; Pfaltz, A. 5-Aza-Semicorrins: A new Class of Bidentate Nitrogen Ligands for Enantioselective Catalysis. *Tetrahedron*, **1992**, *48*, 2143–2156; b) Doyle M. P., Protopopova M. N. New Aspects of Catalytic Asymmetric Cyclopropanation. *Tetrahedron*, **1998**, *54*, 7919–7946; c) Kowalczyk R., Skarzewski J. Selective Nucleophilic Substitution Using Pyrazolate Anions: Easy Approach to Modular Chiral Ligands with Complexing Pyrazole Moieties. *Pol. J. Chem.*, **2007**, *81*, 1987–1999.

[10] **Typical procedure for the synthesis of 5:** A mixture of a bis-imidazole or imidazole ligand **1-4** (0.04 mmol), a drop of hydrazine hydrate, and CuOTf (toluene complex) (0.02 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2 ml), was stirred magnetically at room temperature for 0.5 h. To this solution was added styrene (416 mg, 4 mmol), and stirring was continued for 10 min. From a solution of ethyl diazoacetate (114 mg, 1 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (0.5 ml), prepared in a syringe, a few drops were added in order to start the reaction. The mixture was warmed to 40 °C and stirring was continued at this temperature for 0.5 h, and then cooled to ca. 20 °C. The rest of the ethyl diazoacetate solution was added drop-wise within 4 h, and the mixture was stirred overnight. After removing of the solvent under reduced pressure, the crude product was purified by column chromatography (hexane/AcOEt 95:5). Analytically pure isomers of ethyl 2-phenylcyclopropane-1-carboxylate **5** were obtained as colorless oils and their spectroscopic data fitted well with literature data (*e.g.* ref. [11]). **IR** (KBr):  $\nu$  3087<sub>w</sub>, 3064<sub>w</sub>, 3030<sub>w</sub>, 2981<sub>w</sub>, 2933<sub>w</sub>, 2908<sub>w</sub>, 2872<sub>w</sub>, 1719<sub>s</sub>, 1603<sub>w</sub>, 1496<sub>w</sub>, 1457<sub>m</sub>, 1412<sub>m</sub>, 1336<sub>m</sub>, 1191<sub>s</sub>, 1078<sub>m</sub>, 1040<sub>s</sub>, 1018<sub>m</sub>, 937<sub>m</sub>, 850<sub>w</sub>, 762<sub>s</sub>, 703<sub>s</sub>, 529<sub>w</sub>. **<sup>1</sup>H-NMR** (CDCl<sub>3</sub>): *trans*-**5**:  $\delta$  7.29–7.10 (m, 5H, 5 arom. **H**); 4.18 (q,  $J_{H-H}$  = 7.2, 2H, CH<sub>2</sub>CH<sub>3</sub>); 2.54–2.51 (m, 1H, **HC**(1)); 1.92–1.89 (m, 1H, **HC**(2)); 1.62–1.59 (m, 1H, **HC**(3)); 1.33–1.27 (m, 1H, **HC**(3)); 1.29 (t,  $J_{H-H}$  = 7.2, 3H, CH<sub>2</sub>CH<sub>3</sub>). *cis*-**5**:  $\delta$  7.35–7.08 (m, 5H, 5



arom. **H**); 3.86 (q,  $J_{\text{H-H}} = 7.2$ , 2H, **CH**<sub>2</sub>**CH**<sub>3</sub>); 2.72–2.38 (m, 1H, **HC**(1)); 2.20–1.87 (m, 1H, **HC**(2)); 1.80–1.48 (m, 1H, **HC**(3)); 1.42–1.11 (m, 1H, **HC**(3)); 0.95 (t,  $J_{\text{H-H}} = 7.2$ , 3H, **CH**<sub>2</sub>**CH**<sub>3</sub>). <sup>13</sup>**C**-NMR (CDCl<sub>3</sub>): (*trans*)-**5**:  $\delta$  173.37 (**CO**); 140.12 (1 arom. **C**); 128.48, 126.48, 126.19 (5 arom. **CH**); 60.70 (**CH**<sub>2</sub>**CH**<sub>3</sub>); 26.17 (**C**(2)); 24.17 (**C**(1)); 17.00 (**C**(3)); 14.28 (**CH**<sub>2</sub>**CH**<sub>3</sub>).

[11] a) Chao, D.-J.; Jeon, S.-J.; Kim, H.-S.; Cho, Ch.-S.; Shim, S.-Ch.; Kim, T.-J. Chiral C<sub>2</sub>-Symmetric Bisferrocenyldiamines as Ligands for Transition Metal Catalyzed Asymmetric Cyclopropanation and Aziridination. *Tetrahedron: Asymmetry*, **1999**, *10*, 3833–3848; b) Sanders, Ch. J.; Gillespie, K. M.; Scott, P. Catalyst Structure and the Enantioselective Cyclopropanation of Alkenes by Copper Complexes of Biaryldiimines: the Importance of Ligand Acceleration. *Tetrahedron: Asymmetry*, **2001**, *12*, 1055–1061.

[12] Evans D. A., Woerpel K. A., Scott M. J. Bis(oxazolines) as Ligands for Self-Organizing Chiral Coordination Polymers. Structure of Copper(I) Catalysts for Enantioselective Cyclopropanation of Olefins. *Angew. Chem., Int. Ed. Engl.*, **1992**, *31*, 430–432.